

Radiation from a D-dimensional collision of shock waves: numerical methods

Marco Sampaio
msampaio@ua.pt

Aveiro University & I3N

March 14th, 2013
nr/hep² - IST

Yesterday we have seen:

- How to **formulate the exact initial value problem** for shock wave collision spacetime.
- A **perturbative approximation**, valid **close to the axis at late times**, which is a tower of wave equations with sources.
- Reduced the problem to computation of **scalar functions**

$$F_m^{(n)}(u, v, \rho) = F_{m, \text{Surf}}^{(n)} + F_{m, \text{Vol}}^{(n)}$$

and focused on the surface integrals

- Discussed **numerical** strategy to compute **integration domain**.

Yesterday we have seen:

- How to **formulate the exact initial value problem** for shock wave collision spacetime.
- A **perturbative approximation**, valid **close to the axis at late times**, which is a tower of wave equations with sources.
- Reduced the problem to computation of **scalar functions**

$$F_m^{(n)}(u, v, \rho) = F_{m, \text{Surf}}^{(n)} + F_{m, \text{Vol}}^{(n)}$$

and focused on the surface integrals

- Discussed **numerical** strategy to compute **integration domain**.

Yesterday we have seen:

- How to **formulate the exact initial value problem** for shock wave collision spacetime.
- A **perturbative approximation**, valid **close to the axis at late times**, which is a tower of wave equations with sources.
- Reduced the problem to computation of **scalar functions**

$$F_m^{(n)}(u, v, \rho) = F_{m, \text{Surf}}^{(n)} + F_{m, \text{Vol}}^{(n)}$$

and focused on the surface integrals

- Discussed **numerical** strategy to compute **integration domain**.

Yesterday we have seen:

- How to **formulate the exact initial value problem** for shock wave collision spacetime.
- A **perturbative approximation**, valid **close to the axis at late times**, which is a tower of wave equations with sources.
- Reduced the problem to computation of **scalar functions**

$$F_m^{(n)}(u, v, \rho) = F_{m, \text{Surf}}^{(n)} + F_{m, \text{Vol}}^{(n)}$$

and focused on the surface integrals

- Discussed **numerical** strategy to compute **integration domain**.

Sensitive features of the integrand $I_m^{D,n}(x_*)$

Integrand contains several pieces **to be computed with care**:

- 1 **Polynomials** $Q_m^{D,n}(x_*)$, ... coefficients defined differentially:

Example of task to be done once at the beginning.

(see Mathematica example 2.1)

- 2 D odd contains a potentially **problematic scaling**.

Example where asymptotic expansions can be useful!

(see Mathematica example 2.2)

- 3 **Singularities** @ $x_* = s = \pm 1$, due to

$$\frac{1}{\sqrt{\pm(1-x_*^2)}} = \frac{1}{\sqrt{(1-x_*)(\pm x_* \pm 1)}}$$

- Adapted change of variable $y = y_s^{\text{root}} - kw^2$ & split domains.
- Writing expressions & using series in problematic regions.

(see Mathematica example 2.3)

Sensitive features of the integrand $I_m^{D,n}(x_*)$

Integrand contains several pieces **to be computed with care**:

1 **Polynomials** $Q_m^{D,n}(x_*)$, ... coefficients defined differentially:
Example of task to be done once at the beginning.
(see Mathematica example 2.1)

2 D odd contains a potentially **problematic scaling**.
Example where asymptotic expansions can be useful!
(see Mathematica example 2.2)

3 **Singularities** @ $x_* = s = \pm 1$, due to

$$\frac{1}{\sqrt{\pm(1-x_*^2)}} = \frac{1}{\sqrt{(1-x_*)(\pm x_* \pm 1)}}$$

- Adapted change of variable $y = y_s^{\text{root}} - kw^2$ & split domains.
- Writing expressions & using series in problematic regions.

(see Mathematica example 2.3)

Sensitive features of the integrand $I_m^{D,n}(x_*)$

Integrand contains several pieces **to be computed with care**:

1 Polynomials $Q_m^{D,n}(x_*)$, ... coefficients defined differentially:
Example of task to be done once at the beginning.
(see Mathematica example 2.1)

2 D odd contains a potentially **problematic scaling**.
Example where asymptotic expansions can be useful!
(see Mathematica example 2.2)

3 Singularities @ $x_* = s = \pm 1$, due to

$$\frac{1}{\sqrt{\pm(1-x_*^2)}} = \frac{1}{\sqrt{(1-x_*)(\pm x_* \pm 1)}}$$

- Adapted change of variable $y = y_s^{\text{root}} - kw^2$ & split domains.
- Writing expressions & using series in problematic regions.

(see Mathematica example 2.3)

Sensitive features of the integrand $I_m^{D,n}(x_*)$

Integrand contains several pieces **to be computed with care**:

1 Polynomials $Q_m^{D,n}(x_*)$, ... coefficients defined differentially:
Example of task to be done once at the beginning.
(see Mathematica example 2.1)

2 D odd contains a potentially **problematic scaling**.
Example where asymptotic expansions can be useful!
(see Mathematica example 2.2)

3 **Singularities** @ $x_* = s = \pm 1$, due to

$$\frac{1}{\sqrt{\pm(1-x_*^2)}} = \frac{1}{\sqrt{(1-x_*)(\pm x_* \pm 1)}}$$

- Adapted change of variable $y = y_s^{\text{root}} - kw^2$ & split domains.
- Writing expressions & using series in problematic regions.

(see Mathematica example 2.3)

Plan - Part II

- 1 Radiation extraction
 - First order wave signal
 - Bondi formula vs Pseudo-tensor & extraction
- 2 Higher orders and two dimensional reduction
 - The Volume terms
 - Two dimensional form
 - The 2D Green's functions
- 3 Summary & Conclusions

- 1 Radiation extraction
 - First order wave signal
 - Bondi formula vs Pseudo-tensor & extraction
- 2 Higher orders and two dimensional reduction
 - The Volume terms
 - Two dimensional form
 - The 2D Green's functions
- 3 Summary & Conclusions

Visualisation of the radiation signal $D = 4$

Play Video



- Signal follows **optical rays** & geometrical suppression.
- **Growth** at axis (shorter $\delta\tau$) & far away (pert. breakdown).

Visualisation of the radiation signal $D = 5$

[Play Video](#)

Suppressed tail in D -odd after second optical ray.

Reduced Wave forms



- 1 Radiation extraction**
 - First order wave signal
 - **Bondi formula vs Pseudo-tensor & extraction**

- 2 Higher orders and two dimensional reduction**
 - The Volume terms
 - Two dimensional form
 - The 2D Green's functions

- 3 Summary & Conclusions**

Approximations

D'Eath & Payne have argued that if:

D'Eath and Payne, PRD Volume 46, Number 2, 658, 675 and 694

Approximations

D'Eath & Payne have argued that if:

D'Eath and Payne, PRD Volume 46, Number 2, 658, 675 and 694

- One assumes **analyticity** ($x \equiv \cos \theta$ & using \mathbb{Z}_2 symm.)

$$C(x) \equiv \frac{2}{E_{CM}} \frac{dE}{dx} = \sum_{n=0}^{+\infty} C_n (x^2 - 1)^n$$

Approximations

D'Eath & Payne have argued that if:

D'Eath and Payne, PRD Volume 46, Number 2, 658, 675 and 694

- One assumes **analyticity** ($x \equiv \cos \theta$ & using \mathbb{Z}_2 symm.)

$$C(x) \equiv \frac{2}{E_{CM}} \frac{dE}{dx} = \sum_{n=0}^{+\infty} C_n (x^2 - 1)^n$$

- Then C_n gets **contributions** from perturbations of **order** $< n$.

$$\Rightarrow \text{Inelasticity } \epsilon_{\text{radiated}} = \sum_{n=0}^{+\infty} \frac{C_n (-2)^n n!}{(2n+1)!!}$$

Series **converges** if $\lim_{n \rightarrow +\infty} |C_{n+1}/C_n| \leq 1$.

Approximations

D'Eath & Payne have argued that if:

D'Eath and Payne, PRD Volume 46, Number 2, 658, 675 and 694

- One assumes **analyticity** ($x \equiv \cos \theta$ & using \mathbb{Z}_2 symm.)

$$C(x) \equiv \frac{2}{E_{CM}} \frac{dE}{dx} = \sum_{n=0}^{+\infty} C_n (x^2 - 1)^n$$

- Then C_n gets **contributions** from perturbations of **order** $< n$.

$$\Rightarrow \text{Inelasticity } \epsilon_{\text{radiated}} = \sum_{n=0}^{+\infty} \frac{C_n (-2)^n n!}{(2n+1)!!}$$

Series **converges** if $\lim_{n \rightarrow +\infty} |C_{n+1}/C_n| \leq 1$.

- **Example:**

$D = 4$ we know $C_0 \simeq 0.250$ & $C_0 - 2C_1/3 = 0.163$

$$\Rightarrow C_1/C_0 \simeq 0.52$$

Thus: 1st order \Leftrightarrow isotropic approximation, $n = 0$

2nd order \Leftrightarrow first angular correction, $n = 1$

Approximations

D'Eath & Payne have argued that if:

D'Eath and Payne, PRD Volume 46, Number 2, 658, 675 and 694

- One assumes **analyticity** ($x \equiv \cos \theta$ & using \mathbb{Z}_2 symm.)

$$C(x) \equiv \frac{2}{E_{CM}} \frac{dE}{dx} = \sum_{n=0}^{+\infty} C_n (x^2 - 1)^n$$

- Then C_n gets **contributions** from perturbations of **order** $< n$.

$$\Rightarrow \text{Inelasticity } \epsilon_{\text{radiated}} = \sum_{n=0}^{+\infty} \frac{C_n (-2)^n n!}{(2n+1)!!}$$

Series **converges** if $\lim_{n \rightarrow +\infty} |C_{n+1}/C_n| \leq 1$.

- **Example:**

$D = 4$ we know $C_0 \simeq 0.250$ & $C_0 - 2C_1/3 = 0.163$

$$\Rightarrow C_1/C_0 \simeq 0.52$$

Thus: 1st order \Leftrightarrow isotropic approximation, $n = 0$

2nd order \Leftrightarrow first angular correction, $n = 1$

Radiation extraction (Landau-Lifshitz pseudo-tensor)

- **Radiative** components $h_{ij} = \delta_{ij}H(u, \mathbf{v}, \rho) + \Delta_{ij}(x)E(u, \mathbf{v}, \rho)$
- At first order ($i = 1$) only $E(u, \mathbf{v}, \rho)$
- In our first paper \rightarrow Landau-Lifshitz pseudo-tensor
Yoshino and Shibata, arXiv:0907.2760

$$\begin{aligned} E_{\text{radiated}} &= \int dt \int_{S^{D-2}} \frac{d\text{Energy}}{dS dt} \\ &\simeq \frac{\Omega_{D-3}}{32\pi G_D} \lim_{\hat{\theta} \rightarrow 0, r \rightarrow \infty} \left(r^2 \rho^{D-4} \int h^{ij}{}_{,v} h_{ij,v} dt \right) \\ \frac{E_{\text{radiated}}}{E_{CM}} &\rightarrow \frac{1}{8} \frac{D-2}{D-3} \lim_{\hat{\theta} \rightarrow 0, r \rightarrow \infty} \left(\int (r \rho^{\frac{D-4}{2}} E_{,v})^2 dt \right) \end{aligned}$$

Radiation extraction (Landau-Lifshitz pseudo-tensor)

- **Radiative** components $h_{ij} = \delta_{ij}H(u, \mathbf{v}, \rho) + \Delta_{ij}(x)E(u, \mathbf{v}, \rho)$
- At first order ($i = 1$) only $E(u, \mathbf{v}, \rho)$
- In our first paper \rightarrow Landau-Lifshitz pseudo-tensor
Yoshino and Shibata, arXiv:0907.2760

$$\begin{aligned} E_{\text{radiated}} &= \int dt \int_{S^{D-2}} \frac{d\text{Energy}}{dS dt} \\ &\simeq \frac{\Omega_{D-3}}{32\pi G_D} \lim_{\hat{\theta} \rightarrow 0, r \rightarrow \infty} \left(r^2 \rho^{D-4} \int h^{ij}{}_{,v} h_{ij,v} dt \right) \\ \frac{E_{\text{radiated}}}{E_{CM}} &\rightarrow \frac{1}{8} \frac{D-2}{D-3} \lim_{\hat{\theta} \rightarrow 0, r \rightarrow \infty} \left(\int (r \rho^{\frac{D-4}{2}} E_{,v})^2 dt \right) \end{aligned}$$

Radiation extraction (Landau-Lifshitz pseudo-tensor)

- **Radiative** components $h_{ij} = \delta_{ij}H(u, v, \rho) + \Delta_{ij}(x)E(u, v, \rho)$
- At first order ($i = 1$) only $E(u, v, \rho)$
- In our first paper \rightarrow Landau-Lifshitz pseudo-tensor
Yoshino and Shibata, arXiv:0907.2760

$$\begin{aligned} E_{\text{radiated}} &= \int dt \int_{S^{D-2}} \frac{d\text{Energy}}{dS dt} \\ &\simeq \frac{\Omega_{D-3}}{32\pi G_D} \lim_{\hat{\theta} \rightarrow 0, r \rightarrow \infty} \left(r^2 \rho^{D-4} \int h^{ij}{}_{,v} h_{ij,v} dt \right) \\ \frac{E_{\text{radiated}}}{E_{CM}} &\rightarrow \frac{1}{8} \frac{D-2}{D-3} \lim_{\hat{\theta} \rightarrow 0, r \rightarrow \infty} \left(\int (r \rho^{\frac{D-4}{2}} E_{,v})^2 dt \right) \end{aligned}$$

Radiation extraction (Landau-Lifshitz pseudo-tensor)

- **Radiative** components $h_{ij} = \delta_{ij}H(u, v, \rho) + \Delta_{ij}(x)E(u, v, \rho)$
- At first order ($i = 1$) only $E(u, v, \rho)$
- In our first paper \rightarrow Landau-Lifshitz pseudo-tensor
Yoshino and Shibata, arXiv:0907.2760


$$\begin{aligned} E_{\text{radiated}} &= \int dt \int_{S^{D-2}} \frac{d\text{Energy}}{dS dt} \\ &\simeq \frac{\Omega_{D-3}}{32\pi G_D} \lim_{\hat{\theta} \rightarrow 0, r \rightarrow \infty} \left(r^2 \rho^{D-4} \int h^{ij}{}_{,v} h_{ij,v} dt \right) \\ \frac{E_{\text{radiated}}}{E_{CM}} &\rightarrow \frac{1}{8} \frac{D-2}{D-3} \lim_{\hat{\theta} \rightarrow 0, r \rightarrow \infty} \left(\int (r \rho^{\frac{D-4}{2}} E_{,v})^2 dt \right) \end{aligned}$$

Radiation extraction (Landau-Lifshitz pseudo-tensor)


- **Radiative** components $h_{ij} = \delta_{ij}H(u, v, \rho) + \Delta_{ij}(x)E(u, v, \rho)$
- At first order ($i = 1$) only $E(u, v, \rho)$
- In our first paper \rightarrow Landau-Lifshitz pseudo-tensor
Yoshino and Shibata, arXiv:0907.2760

$$\begin{aligned} E_{\text{radiated}} &= \int dt \int_{S^{D-2}} \frac{d\text{Energy}}{dS dt} \\ &\simeq \frac{\Omega_{D-3}}{32\pi G_D} \lim_{\hat{\theta} \rightarrow 0, r \rightarrow \infty} \left(r^2 \rho^{D-4} \int h^{ij}{}_{,v} h_{ij,v} dt \right) \\ \frac{E_{\text{radiated}}}{E_{CM}} &\rightarrow \frac{1}{8} \frac{D-2}{D-3} \lim_{\hat{\theta} \rightarrow 0, r \rightarrow \infty} \left(\int (r \rho^{\frac{D-4}{2}} E_{,v})^2 dt \right) \end{aligned}$$


Bondi extraction & Higher orders ...



Bondi extraction & Higher orders ...




Bondi extraction & Higher orders ...




Bondi extraction & Higher orders ...

Tanabe, Kinoshita and Shiromizu, 1104.0303



Bondi extraction & Higher orders ...

Tanabe, Kinoshita and Shiromizu, 1104.0303



de Donder coordinates \rightarrow **Bondi** coordinates:


$$ds^2 = g_{\hat{\tau}\hat{\tau}} d\hat{\tau}^2 + 2g_{\hat{\tau}\hat{r}} d\hat{\tau} d\hat{r} + 2g_{\hat{\imath}\hat{\tau}} dx^{\hat{\imath}} d\hat{\tau} + \hat{r}^2 \left[\omega_{\hat{\imath}\hat{\jmath}} + \frac{h_{\hat{\imath}\hat{\jmath}}^{[1]}}{\hat{r}^{D/2-1}} + \dots \right] dx^{\hat{\imath}} dx^{\hat{\jmath}}$$

■ Conditions: $g_{\hat{r}\hat{r}} = g_{\hat{r}\hat{\theta}} = 0$ & \hat{r} areal radius

■ Use de Donder gauge conditions: $\bar{h}^{\alpha\beta}_{,\beta} = 0$

Bondi extraction & Higher orders ...

Tanabe, Kinoshita and Shiromizu, 1104.0303



de Donder coordinates \rightarrow **Bondi** coordinates:



$$ds^2 = g_{\hat{\tau}\hat{\tau}} d\hat{\tau}^2 + 2g_{\hat{\tau}\hat{r}} d\hat{\tau} d\hat{r} + 2g_{\hat{\imath}\hat{\jmath}} dx^{\hat{\imath}} d\hat{\tau} + \hat{r}^2 \left[\omega_{\hat{\imath}\hat{\jmath}} + \frac{\dot{h}_{\hat{\imath}\hat{\jmath}}^{[1]}}{\hat{r}^{D/2-1}} + \dots \right] dx^{\hat{\imath}} dx^{\hat{\jmath}}$$

■ Conditions: $g_{\hat{r}\hat{r}} = g_{\hat{r}\hat{\theta}} = 0$ & \hat{r} areal radius

■ Use de Donder gauge conditions: $\bar{h}^{\alpha\beta}_{,\beta} = 0$


$$\Rightarrow \frac{dM_B}{d\hat{\tau} d \cos \hat{\theta}} = -\frac{(D-2)(D-3)\Omega_{D-3}}{32\pi G_D} \lim_{\hat{r} \rightarrow +\infty} \left[\hat{r} \hat{\rho}^{\frac{D-4}{2}} (\dot{E} + \dot{H}) \right]^2$$

Extracting the inelasticity @ 1st order




D	4	5	6	7	8	9	10
$\epsilon_{1st\ order}(\%)$	25.0	30.0	33.3	35.7	37.5	38.9	40.0
AH bound (%)	29.3	33.5	36.1	37.9	39.3	40.4	41.2

A remarkably simple fit formula



A remarkably simple fit formula



$$\epsilon_{\text{1st order}} = \frac{1}{2} - \frac{1}{D} \quad (!!!)$$

Outline

- 1 Radiation extraction
 - First order wave signal
 - Bondi formula vs Pseudo-tensor & extraction
- 2 Higher orders and two dimensional reduction
 - The Volume terms
 - Two dimensional form
 - The 2D Green's functions
- 3 Summary & Conclusions


Volume terms & domain conditions

Recall:

$$F_{m,Vol}^{(n)} = \frac{-\Omega_{D-4}}{2(2\pi\rho)^{\frac{D-2}{2}}} \int_0^u du' \iint_{\mathcal{D}_{Vol}} dv' d\rho' \rho'^{\frac{D-4}{2}} S_F^{(n-1)}(u', v', \rho') I_m^{D,0}(x)$$

$$x = \frac{\rho^2 + \rho'^2 - 2(u-u')(v-v')}{2\rho\rho'} \leq 1$$

($I_m^{D,0}$ in handout).




Volume terms & domain conditions

Recall:

$$F_{m,Vol}^{(n)} = \frac{-\Omega_{D-4}}{2(2\pi\rho)^{\frac{D-2}{2}}} \int_0^u du' \iint_{\mathcal{D}_{Vol}} dv' d\rho' \rho'^{\frac{D-4}{2}} S_F^{(n-1)}(u', v', \rho') I_m^{D,0}(x)$$

$$x = \frac{\rho^2 + \rho'^2 - 2(u-u')(v-v')}{2\rho\rho'} \leq 1$$

($I_m^{D,0}$ in handout).



This requires:

- Evaluating ~ 10 surface terms in $S_F^{(n-1)}$
- Triple integration
- Finding a complicated implicit integration domain in 2D
(Also need to impose above blue surface)


Volume terms & domain conditions

Recall:

$$F_{m,Vol}^{(n)} = \frac{-\Omega_{D-4}}{2(2\pi\rho)^{\frac{D-2}{2}}} \int_0^u du' \iint_{\mathcal{D}_{Vol}} dv' d\rho' \rho'^{\frac{D-4}{2}} S_F^{(n-1)}(u', v', \rho') I_m^{D,0}(x)$$

$$x = \frac{\rho^2 + \rho'^2 - 2(u-u')(v-v')}{2\rho\rho'} \leq 1$$

($I_m^{D,0}$ in handout).



This requires:

- Evaluating ~ 10 surface terms in $S_F^{(n-1)}$
- **Triple** integration
- Finding a **complicated** implicit integration **domain in 2D**
(Also need to impose above blue surface)


Volume terms & domain conditions

Recall:

$$F_{m,Vol}^{(n)} = \frac{-\Omega_{D-4}}{2(2\pi\rho)^{\frac{D-2}{2}}} \int_0^u du' \iint_{\mathcal{D}_{Vol}} dv' d\rho' \rho'^{\frac{D-4}{2}} S_F^{(n-1)}(u', v', \rho') I_m^{D,0}(x)$$


$$x = \frac{\rho^2 + \rho'^2 - 2(u-u')(v-v')}{2\rho\rho'} \leq 1$$

($I_m^{D,0}$ in handout).



This requires:

- Evaluating ~ 10 surface terms in $S_F^{(n-1)}$
- Triple integration
- Finding a complicated implicit integration domain in 2D
(Also need to impose above blue surface)



Source functions

Most important sources are in ij radiative components:

$$T_{ij}^{(1)} = T^{(1)}(u, v, \rho)\delta_{ij} + S^{(1)}(u, v, \rho)\Delta_{ij}$$

Source functions

Most important sources are in ij radiative components:

$$T_{ij}^{(1)} = T^{(1)}(u, v, \rho)\delta_{ij} + S^{(1)}(u, v, \rho)\Delta_{ij}$$

and after a lengthy calculation (see lecture notes)

$$T^{(1)} = -\frac{1}{D-2}(B_{,v})^2 - \frac{2(D-3)}{D-2} \left[\frac{D-3}{\rho} BE_{,v} + B_{,\rho} E_{,v} + BE_{,\rho v} \right] - 2(D-3)E_{,u}E_{,v} + \\ + (D-3) \left[\frac{4(D-3)}{\rho^2} E^2 + \frac{D}{2(D-2)} (E_{,\rho})^2 - \frac{D^2-9D+16}{(D-2)\rho} EE_{,\rho} - \frac{D-4}{D-2} EE_{,\rho\rho} \right]$$

Source functions

Most important sources are in ij radiative components:

$$T_{ij}^{(1)} = T^{(1)}(u, v, \rho)\delta_{ij} + S^{(1)}(u, v, \rho)\Delta_{ij}$$

and after a lengthy calculation (see lecture notes)

$$T^{(1)} = -\frac{1}{D-2}(B_{,v})^2 - \frac{2(D-3)}{D-2} \left[\frac{D-3}{\rho} BE_{,v} + B_{,\rho} E_{,v} + BE_{,\rho v} \right] - 2(D-3)E_{,u}E_{,v} + \\ + (D-3) \left[\frac{4(D-3)}{\rho^2} E^2 + \frac{D}{2(D-2)}(E_{,\rho})^2 - \frac{D^2-9D+16}{(D-2)\rho} EE_{,\rho} - \frac{D-4}{D-2} EE_{,\rho\rho} \right]$$

and

$$S^{(1)} = AE_{,vv} + \frac{1}{D-2} \left[(B_{,v})^2 - \frac{2(D-3)}{\rho} BE_{,v} + 2(D-3)B_{,\rho} E_{,v} - 2BE_{,\rho v} \right] + \\ + \frac{4(D-3)}{\rho^2} E^2 - \frac{2(D-3)^2}{(D-2)\rho} EE_{,\rho} - \frac{2(D-3)}{D-2} EE_{,\rho\rho} + (D-4) \left[2E_{,u}E_{,v} - \frac{D-1}{2(D-2)}(E_{,\rho})^2 \right]$$

Source functions

Most important sources are in ij radiative components:

$$T_{ij}^{(1)} = T^{(1)}(u, v, \rho)\delta_{ij} + S^{(1)}(u, v, \rho)\Delta_{ij}$$

and after a lengthy calculation (see lecture notes)

$$T^{(1)} = -\frac{1}{D-2}(B_{,v})^2 - \frac{2(D-3)}{D-2} \left[\frac{D-3}{\rho} BE_{,v} + B_{,\rho} E_{,v} + BE_{,\rho v} \right] - 2(D-3)E_{,u}E_{,v} + \\ + (D-3) \left[\frac{4(D-3)}{\rho^2} E^2 + \frac{D}{2(D-2)}(E_{,\rho})^2 - \frac{D^2-9D+16}{(D-2)\rho} EE_{,\rho} - \frac{D-4}{D-2} EE_{,\rho\rho} \right]$$

and

$$S^{(1)} = AE_{,vv} + \frac{1}{D-2} \left[(B_{,v})^2 - \frac{2(D-3)}{\rho} BE_{,v} + 2(D-3)B_{,\rho} E_{,v} - 2BE_{,\rho v} \right] + \\ + \frac{4(D-3)}{\rho^2} E^2 - \frac{2(D-3)^2}{(D-2)\rho} EE_{,\rho} - \frac{2(D-3)}{D-2} EE_{,\rho\rho} + (D-4) \left[2E_{,u}E_{,v} - \frac{D-1}{2(D-2)}(E_{,\rho})^2 \right]$$

\Rightarrow Some simplification is in order for feasibility!

Outline

- 1 Radiation extraction
 - First order wave signal
 - Bondi formula vs Pseudo-tensor & extraction
- 2 Higher orders and two dimensional reduction
 - The Volume terms
 - **Two dimensional form**
 - The 2D Green's functions
- 3 Summary & Conclusions

2D reduction & CL symmetry

D'Eath & Payne found conformal symmetry (valid in $D > 4$)

2D reduction & CL symmetry

D'Eath & Payne found conformal symmetry (valid in $D > 4$)

For one shock it consists of a one-parameter symmetry:

- **Boost:** Scales up energy parameter $E \rightarrow E' = e^\beta E$ preserving transverse coordinates.
- **+ Conformal scaling:** as to bring E' back to E .

2D reduction & CL symmetry

D'Eath & Payne found conformal symmetry (valid in $D > 4$)

For one shock it consists of a one-parameter symmetry:

- **Boost:** Scales up energy parameter $E \rightarrow E' = e^\beta E$ preserving transverse coordinates.
- **+ Conformal scaling:** as to bring E' back to E .

This also **holds** for system of shocks **order by order**.

Thus:

- One finds $D - 1$ invariant coordinates $\{p, q, \phi_i\}$
- All ρ dependence scales out!
- \Rightarrow Problem becomes 1+1 in $\{p, q\}$ order by order

(Recall $p \equiv (\sqrt{2}v - \Phi(\rho))\rho^{D-4}$ and $q \equiv \frac{u}{\rho^{D-2}}$)

Integral solutions in 2D form

More specifically, the symmetry implies:

$$h_{\mu\nu}^{(i)}(\mathbf{p}, \mathbf{q}, \phi_i, \rho) = \rho^{-(D-3)(2i+N_u-N_v)} f_{\mu\nu}^{(i)}(\mathbf{p}, \mathbf{q}, \phi_i) .$$

Integral solutions in 2D form

More specifically, the symmetry implies:

$$h_{\mu\nu}^{(i)}(\rho, q, \phi_i, \rho) = \rho^{-(D-3)(2i+N_u-N_v)} f_{\mu\nu}^{(i)}(\rho, q, \phi_i) .$$

And so the previous solutions acquire the form

$$f_{m, \text{Surf}}^{(n)}(\rho, q) = -\frac{n!(-1)^D \Omega_{D-4}}{(2\pi)^{\frac{D-2}{2}}} \left(\frac{\sqrt{2}}{q}\right)^n \int_{\mathcal{D}_{\text{surf}}} dy f(y) y^{\frac{D-4}{2}+n} I_m^{D,n}(x_*)$$

Integral solutions in 2D form

More specifically, the symmetry implies:

$$h_{\mu\nu}^{(i)}(p, q, \phi_i, \rho) = \rho^{-(D-3)(2i+N_u-N_v)} f_{\mu\nu}^{(i)}(p, q, \phi_i) .$$

And so the previous solutions acquire the form

$$f_{m, \text{Surf}}^{(n)}(p, q) = -\frac{n!(-1)^D \Omega_{D-4}}{(2\pi)^{\frac{D-2}{2}}} \left(\frac{\sqrt{2}}{q}\right)^n \int_{\mathcal{D}_{\text{surf}}} dy f(y) y^{\frac{D-4}{2}+n} I_m^{D,n}(x_*)$$

and

$$f_{m, \text{Vol}}^{(n)}(p, q) = \frac{(-1)^D}{2} \iint dq' dp' \sigma^{(n-1)}(p', q') G^{(n,m)}(p, q, p', q')$$

Integral solutions in 2D form

More specifically, the symmetry implies:

$$h_{\mu\nu}^{(i)}(p, q, \phi_i, \rho) = \rho^{-(D-3)(2i+N_u-N_v)} f_{\mu\nu}^{(i)}(p, q, \phi_i).$$

And so the previous solutions acquire the form

$$f_{m, \text{Surf}}^{(n)}(p, q) = -\frac{n!(-1)^D \Omega_{D-4}}{(2\pi)^{\frac{D-2}{2}}} \left(\frac{\sqrt{2}}{q}\right)^n \int_{\mathcal{D}_{\text{surf}}} dy f(y) y^{\frac{D-4}{2}+n} I_m^{D,n}(x_*)$$

and

$$f_{m, \text{Vol}}^{(n)}(p, q) = \frac{(-1)^D}{2} \iint dq' dp' \sigma^{(n-1)}(p', q') G^{(n,m)}(p, q, p', q')$$


with

$$G^{(n,m)}(p, q, p', q') \equiv \frac{(-1)^{D+1} \Omega_{D-4}}{(2\pi)^{\frac{D-2}{2}}} \int_{\mathcal{D}'_{\text{surf}}} dy y^{\frac{D-4}{2}-2n(D-3)} I_m^{D,0}(x)$$


$$x = \frac{1+y^2-\sqrt{2}(q-q'y^{D-2})(p-p'y^{-(D-4)}-\psi(y))}{2y} \leq 1$$

$$\Rightarrow \hat{C}_-(y) \leq 0 \text{ (Note: When } p' = q' = 0, \hat{C}_-(y) \rightarrow C_-(y))$$


First order results in 2D form




First order results in 2D form




First order results in 2D form





First order results in 2D form






First order results in 2D form







First order results in 2D form



First order results in 2D form





First order results in 2D form



Outline

- 1 Radiation extraction
 - First order wave signal
 - Bondi formula vs Pseudo-tensor & extraction
- 2 Higher orders and two dimensional reduction
 - The Volume terms
 - Two dimensional form
 - The 2D Green's functions
- 3 Summary & Conclusions

Characteristics & Past light cone visualisation



Domain conditions & generalised domain search

In 2D form the **domain condition** (defining light-cone) is a more **complicated** polynomial. **The procedure becomes:**

- Construct a chain of derivatives $C_s^{(k)} \sim \frac{d\hat{C}_s(y)}{dy}$, $k = 0, \dots, 3$
- Find root of $C_s^{(3)}$ and use that to find possible root configuration for $C_s^{(2)}$.
- Proceed **iteratively** down to \hat{C}_s .

To perform the source (double) integral in the 2D plane, need generalised domain search \Rightarrow **2D bracketing strategy**.

Domain conditions & generalised domain search

In 2D form the **domain condition** (defining light-cone) is a more **complicated** polynomial. **The procedure becomes:**

- Construct a chain of derivatives $C_s^{(k)} \sim \frac{d\hat{C}_s(y)}{dy}$, $k = 0, \dots, 3$
- Find root of $C_s^{(3)}$ and use that to find possible root configuration for $C_s^{(2)}$.
- Proceed **iteratively** down to \hat{C}_s .

To perform the source (double) integral in the 2D plane, need generalised domain search \Rightarrow **2D bracketing strategy**.

Domain conditions & generalised domain search

In 2D form the **domain condition** (defining light-cone) is a more **complicated** polynomial. **The procedure becomes:**

- Construct a chain of derivatives $C_s^{(k)} \sim \frac{d\hat{C}_s(y)}{dy}$, $k = 0, \dots, 3$
- Find root of $C_s^{(3)}$ and use that to find possible root configuration for $C_s^{(2)}$.
- Proceed **iteratively** down to \hat{C}_s .

To perform the source (double) integral in the 2D plane, need generalised domain search \Rightarrow **2D bracketing strategy**.

Domain conditions & generalised domain search

In 2D form the **domain condition** (defining light-cone) is a more **complicated** polynomial. **The procedure becomes:**

- Construct a chain of derivatives $C_s^{(k)} \sim \frac{d\hat{C}_s(y)}{dy}$, $k = 0, \dots, 3$
- Find root of $C_s^{(3)}$ and use that to find possible root configuration for $C_s^{(2)}$.
- Proceed **iteratively** down to \hat{C}_s .

To perform the source (double) integral in the 2D plane, need generalised domain search \Rightarrow **2D bracketing strategy**.

Collisions of AS shock waves \Rightarrow interesting to study:

- **Inelasticity of BH formation** in transplanckian collisions.
- **Perturbative techniques** in the context of strong gravity.
- **Numerical strategies** to solve higher dimensional integrals in non-trivial domains.

The results so far:

- Show **consistency** with the apparent horizon bound.
- A **puzzling simplicity** in first order perturbation theory.

Collisions of AS shock waves \Rightarrow interesting to study:

- **Inelasticity of BH formation** in transplanckian collisions.
- **Perturbative techniques** in the context of strong gravity.
- **Numerical strategies** to solve higher dimensional integrals in non-trivial domains.

The results so far:

- Show **consistency** with the apparent horizon bound.
- A **puzzling simplicity** in first order perturbation theory.

Collisions of AS shock waves \Rightarrow interesting to study:

- **Inelasticity of BH formation** in transplanckian collisions.
- **Perturbative techniques** in the context of strong gravity.
- **Numerical strategies** to solve higher dimensional integrals in non-trivial domains.

The results so far:

- Show **consistency** with the apparent horizon bound.
- A **puzzling simplicity** in first order perturbation theory.

Collisions of AS shock waves \Rightarrow interesting to study:

- **Inelasticity of BH formation** in transplanckian collisions.
- **Perturbative techniques** in the context of strong gravity.
- **Numerical strategies** to solve higher dimensional integrals in non-trivial domains.

The results so far:

- Show **consistency** with the apparent horizon bound.
- A **puzzling simplicity** in first order perturbation theory.

Thanks for your attention!