

Radiation from a D-dimensional collision of shock waves: numerical methods

Marco Sampaio
msampaio@ua.pt

Aveiro University & I3N

March 13th, 2013
nr/hep² - IST

Acknowledgments

■ Collaborators

Flávio Coelho, Carlos Herdeiro & Carmen Rebelo

JHEP07(2011)121

PRL 108 (2012) 181102

arXiv:1206.5839

■ Funding & Institutions



universidade de aveiro

Gr@v

gravitation.web.ua.pt





■ nr/hep² organizers

Motivation I – The Transplanckian Problem

Motivation I – The Transplanckian Problem

- 1 Large extra dimensions (ADD) address **hierarchy problem**
Transplanckian scattering @ $\sim 1\text{TeV} \Rightarrow$ **BHs @ LHC**



$v/c > 0.999$ @ LHC

CMS arXiv:1012.3357



ATLAS-CONF-2011-065

rough model Gr. Rad.

Frost, MS, Gaunt et. al. 0904.0979

Motivation I – The Transplanckian Problem

- 1 Large extra dimensions (ADD) address **hierarchy problem**
Transplanckian scattering @ $\sim 1\text{TeV} \Rightarrow$ **BHs @ LHC**



$v/c > 0.999$ @ LHC

CMS arXiv:1012.3357

ATLAS-CONF-2011-065

rough model Gr. Rad.

Frost, MS, Gaunt et. al. 0904.0979


- 2 Numerical **GR**, BH collisions difficult for **large boost**.

U. Sperhake, V. Cardoso, F. Pretorius, E. Berti, J. Gonzalez, arXiv:0806.1738

M. Shibata, H. Okawa, T. Yamamoto, arXiv:0810.4735 + etc ...

Motivation I – The Transplanckian Problem

- 1 Large extra dimensions (ADD) address **hierarchy problem**
Transplanckian scattering @ $\sim 1\text{TeV} \Rightarrow$ **BHs @ LHC**



$v/c > 0.999$ @ LHC

CMS arXiv:1012.3357

ATLAS-CONF-2011-065

rough model Gr. Rad.

Frost, MS, Gaunt et. al. 0904.0979

- 2 Numerical **GR**, BH collisions difficult for **large boost**.

U. Sperhake, V. Cardoso, F. Pretorius, E. Berti, J. Gonzalez, arXiv:0806.1738

M. Shibata, H. Okawa, T. Yamamoto, arXiv:0810.4735 + etc ...

- 3 Shock wave collisions in $D \geq 4$:


- In $D = 4$, $\epsilon_{\text{rad}}^{(1)} \simeq 0.25$, & $\epsilon_{\text{rad}}^{(2)} \simeq 0.163$ (**within Num. GR err**)

D'Eath and Payne, PRD Volume 46, Number 2, 658, 675 and 694

see also East and Pretorius, arXiv:1210.0443

Motivation I – The Transplanckian Problem

- 1 Large extra dimensions (ADD) address **hierarchy problem**
Transplanckian scattering @ $\sim 1\text{TeV} \Rightarrow$ **BHs @ LHC**



$v/c > 0.999$ @ LHC

CMS arXiv:1012.3357

ATLAS-CONF-2011-065

rough model Gr. Rad.

Frost, MS, Gaunt et. al. 0904.0979

- 2 Numerical **GR**, BH collisions difficult for **large boost**.

U. Sperhake, V. Cardoso, F. Pretorius, E. Berti, J. Gonzalez, arXiv:0806.1738

M. Shibata, H. Okawa, T. Yamamoto, arXiv:0810.4735 + etc ...

- 3 Shock wave collisions in $D \geq 4$:

- In $D = 4$, $\epsilon_{\text{rad}}^{(1)} \simeq 0.25$, & $\epsilon_{\text{rad}}^{(2)} \simeq 0.163$ (**within Num. GR err**)

D'Eath and Payne, PRD Volume 46, Number 2, 658, 675 and 694

see also East and Pretorius, arXiv:1210.0443

- In $D \geq 4$, $\epsilon_{\text{rad}}^{(1)} = ?$, $\epsilon_{\text{rad}}^{(2)} = ?$

Motivation II – Numerics vs Analytics

This problem is also technically very rich

1 Analytically:

- Allows a consistent setup of an evolution problem with well defined *initial conditions*,
- An *approximate integral solution* can be obtained using *perturbation theory*,
- Requires the discussion of a general radiation extraction formula in axially symmetric spacetimes at null infinity.

2 Numerically it requires:

- The integration of many non-trivial integral solutions using Green functions,
- The discussion of strategies to compute non-trivial integration domains,
- The treatment of singularities, and other numerical issues through series expansions & recurrence relations.

Motivation II – Numerics vs Analytics

This problem is also technically very rich

1 Analytically:

- Allows a **consistent setup of an evolution problem** with well defined *initial conditions*,
- An *approximate integral solution* can be obtained using **perturbation theory**,
- Requires the discussion of a general **radiation extraction** formula in axially symmetric spacetimes **at null infinity**.

2 Numerically it requires:

- The **integration** of many non-trivial integral **solutions using Green functions**,
- The discussion of strategies to **compute non-trivial integration domains**,
- The treatment of **singularities**, and other numerical issues through **series expansions & recurrence relations**.

Motivation II – Numerics vs Analytics

This problem is also technically very rich

1 Analytically:

- Allows a **consistent setup of an evolution problem** with well defined *initial conditions*,
- An *approximate integral solution* can be obtained using **perturbation theory**,
- Requires the discussion of a general **radiation extraction** formula in axially symmetric spacetimes **at null infinity**.

2 Numerically it requires:

- The **integration** of many non-trivial integral **solutions using Green functions**,
- The discussion of strategies to **compute non-trivial integration domains**,
- The treatment of **singularities**, and other numerical issues through **series expansions & recurrence relations**.

Motivation II – Numerics vs Analytics

This problem is also technically very rich

1 Analytically:

- Allows a **consistent setup of an evolution problem** with well defined *initial conditions*,
- An *approximate integral solution* can be obtained using **perturbation theory**,
- Requires the discussion of a general **radiation extraction** formula in axially symmetric spacetimes **at null infinity**.

2 Numerically it requires:

- The **integration** of many non-trivial integral **solutions using Green functions**,
- The discussion of strategies to **compute non-trivial integration domains**,
- The treatment of **singularities**, and other numerical issues through **series expansions & recurrence relations**.

Motivation II – Numerics vs Analytics

This problem is also technically very rich

1 Analytically:

- Allows a **consistent setup of an evolution problem** with well defined *initial conditions*,
- An *approximate integral solution* can be obtained using **perturbation theory**,
- Requires the discussion of a general **radiation extraction** formula in axially symmetric spacetimes **at null infinity**.

2 Numerically it requires:

- The **integration** of many non-trivial integral **solutions using Green functions**,
- The discussion of strategies to **compute non-trivial integration domains**,
- The treatment of **singularities**, and other numerical issues through **series expansions & recurrence relations**.

Motivation II – Numerics vs Analytics

This problem is also technically very rich

1 Analytically:

- Allows a **consistent setup of an evolution problem** with well defined *initial conditions*,
- An *approximate integral solution* can be obtained using **perturbation theory**,
- Requires the discussion of a general **radiation extraction** formula in axially symmetric spacetimes **at null infinity**.

2 Numerically it requires:

- The **integration** of many non-trivial integral **solutions using Green functions**,
- The discussion of strategies to **compute non-trivial integration domains**,
- The treatment of **singularities**, and other numerical issues through **series expansions & recurrence relations**.

Motivation II – Numerics vs Analytics

This problem is also technically very rich

1 Analytically:

- Allows a **consistent setup of an evolution problem** with well defined *initial conditions*,
- An *approximate integral solution* can be obtained using **perturbation theory**,
- Requires the discussion of a general **radiation extraction** formula in axially symmetric spacetimes **at null infinity**.

2 Numerically it requires:

- The **integration** of many non-trivial integral **solutions using Green functions**,
- The discussion of strategies to **compute non-trivial integration domains**,
- The treatment of **singularities**, and other numerical issues through **series expansions & recurrence relations**.

Motivation II – Numerics vs Analytics

This problem is also technically very rich

1 Analytically:

- Allows a **consistent setup of an evolution problem** with well defined *initial conditions*,
- An *approximate integral solution* can be obtained using **perturbation theory**,
- Requires the discussion of a general **radiation extraction** formula in axially symmetric spacetimes **at null infinity**.

2 Numerically it requires:

- The **integration** of many non-trivial integral **solutions using Green functions**,
- The discussion of strategies to **compute non-trivial integration domains**,
- The treatment of **singularities**, and other numerical issues through **series expansions & recurrence relations**.

⇒ Excelent ground to illustrate **many** techniques, within a well defined and **interesting problem**.

- 1 AS Gravitational Shock Waves
 - Physical properties & Ray Optics
 - Superposition & causal structure
 - Perturbative Setup & Solutions

- 2 Numerical Strategies – Surface Integrals
 - Breaking down the problem
 - Numerical domain search
 - Dealing with the Integrand

Plan for Today

- 1 AS Gravitational Shock Waves
 - Physical properties & Ray Optics
 - Superposition & causal structure
 - Perturbative Setup & Solutions

- 2 Numerical Strategies – Surface Integrals
 - Breaking down the problem
 - Numerical domain search
 - Dealing with the Integrand

Tomorrow: Radiation extraction, Sources & Results.

- 1 AS Gravitational Shock Waves**
 - Physical properties & Ray Optics
 - Superposition & causal structure
 - Perturbative Setup & Solutions

- 2 Numerical Strategies – Surface Integrals**
 - Breaking down the problem
 - Numerical domain search
 - Dealing with the Integrand

The Aichelburg-Sexl ultraboost


The Aichelburg-Sexl ultraboost

Schwarzschild metric \rightarrow field of source $\mu \propto G_D M$ at rest.

The Aichelburg-Sexl ultraboost

Schwarzschild metric \rightarrow field of source $\mu \propto G_D M$ at rest.


$$ds^2 = - \left(1 - \frac{\mu}{r^{D-3}}\right) dt^2 + \left(1 - \frac{\mu}{r^{D-3}}\right)^{-1} dr^2 + r^2 d\Omega_{D-2}^2$$



The Aichelburg-Sexl ultraboost

Schwarzschild metric \rightarrow field of source $\mu \propto G_D M$ at rest.


$$ds^2 = - \left(1 - \frac{\mu}{r^{D-3}}\right) dt^2 + \left(1 - \frac{\mu}{r^{D-3}}\right)^{-1} dr^2 + r^2 d\Omega_{D-2}^2$$



The Aichelburg-Sexl ultraboost

Schwarzschild metric \rightarrow field of source $\mu \propto G_D M$ at rest.


$$ds^2 = - \left(1 - \frac{\mu}{r^{D-3}}\right) dt^2 + \left(1 - \frac{\mu}{r^{D-3}}\right)^{-1} dr^2 + r^2 d\Omega_{D-2}^2$$




The Aichelburg-Sexl ultraboost

Schwarzschild metric \rightarrow field of source $\mu \propto G_D M$ at rest.

$$ds^2 = - \left(1 - \frac{\mu}{r^{D-3}}\right) dt^2 + \left(1 - \frac{\mu}{r^{D-3}}\right)^{-1} dr^2 + r^2 d\Omega_{D-2}^2$$



$$ds^2 = - dudv + d\rho^2 + \rho^2 d\Omega_{D-3}^2 + \kappa\Phi(\rho)\delta(u)du^2$$




$$(u, v) = (t - z, t + z)$$

The Aichelburg-Sexl ultraboost


Schwarzschild metric \rightarrow field of source $\mu \propto G_D M$ at rest.

$$ds^2 = - \left(1 - \frac{\mu}{r^{D-3}}\right) dt^2 + \left(1 - \frac{\mu}{r^{D-3}}\right)^{-1} dr^2 + r^2 d\Omega_{D-2}^2$$



$$ds^2 = - dudv + d\rho^2 + \rho^2 d\Omega_{D-3}^2 + \kappa\Phi(\rho)\delta(u)du^2$$

Flat region II



Flat region I

$$(u, v) = (t - z, t + z)$$

Basic properties of a single shock wave I

- Solution of Einstein's equations, **point source** $P^\mu = E n^\mu$

$$T^{\mu\nu} = E \delta(u) \delta^{(D-2)}(x^i) n^\mu n^\nu, \quad n^\mu n_\mu = 0, \quad \kappa = \frac{8\pi G_D}{\Omega_{D-3}} E$$


- The **shock profile** is $\Phi(\rho) = \begin{cases} \frac{2}{(D-4)\rho^{D-4}}, & D > 4 \end{cases}$.

Basic properties of a single shock wave I

- Solution of Einstein's equations, **point source** $P^\mu = E n^\mu$

$$T^{\mu\nu} = E \delta(u) \delta^{(D-2)}(x^i) n^\mu n^\nu, \quad n^\mu n_\mu = 0, \quad \kappa = \frac{8\pi G_D}{\Omega_{D-3}} E$$

- The **shock profile** is $\Phi(\rho) = \begin{cases} \frac{2}{(D-4)\rho^{D-4}}, & D > 4 \end{cases}$.




Basic properties of a single shock wave I

- Solution of Einstein's equations, **point source** $P^\mu = E n^\mu$

$$T^{\mu\nu} = E \delta(u) \delta^{(D-2)}(x^i) n^\mu n^\nu, \quad n^\mu n_\mu = 0, \quad \kappa = \frac{8\pi G_D}{\Omega_{D-3}} E$$

- The **shock profile** is $\Phi(\rho) = \begin{cases} \frac{2}{(D-4)\rho^{D-4}}, & D > 4 \\ -2 \ln(\rho), & D = 4 \end{cases}$.



Basic properties of a single shock wave II

Symmetries:

- **Axial symmetry** (ϕ_i rotations on $d\Omega_{D-3}$)
- Advanced **null translations** $v \rightarrow v + \text{const.}$
 \Rightarrow Metric of **one shock** wave effectively **2D** (u, ρ)
- **Boosts** along $+z$, with velocity $\beta \equiv \tanh \alpha$, **up to a scaling**

$$E \rightarrow E' = e^\alpha E \quad (\Leftrightarrow \kappa \rightarrow \nu = e^\alpha E)$$

Geometrical optics:

- **Riemann** tensor **singular** on the shock plane
- **Null** rays & tangent vectors **discontinuous** across shock

Basic properties of a single shock wave II

Symmetries:

- **Axial symmetry** (ϕ_i rotations on $d\Omega_{D-3}$)
- Advanced **null translations** $v \rightarrow v + \text{const.}$
 \Rightarrow Metric of **one shock** wave effectively **2D** (u, ρ)
- **Boosts** along $+z$, with velocity $\beta \equiv \tanh \alpha$, **up to a scaling**

$$E \rightarrow E' = e^\alpha E \quad (\Leftrightarrow \kappa \rightarrow \nu = e^\alpha E)$$

Geometrical optics:

- **Riemann** tensor **singular** on the shock plane
- **Null** rays & tangent vectors **discontinuous** across shock

Basic properties of a single shock wave II

Symmetries:

- **Axial symmetry** (ϕ_i rotations on $d\Omega_{D-3}$)
- Advanced **null translations** $v \rightarrow v + \text{const.}$
 \Rightarrow Metric of **one shock** wave effectively **2D** (u, ρ)
- **Boosts** along $+z$, with velocity $\beta \equiv \tanh \alpha$, **up to a scaling**

$$E \rightarrow E' = e^\alpha E \quad (\Leftrightarrow \kappa \rightarrow \nu = e^\alpha E)$$

Geometrical optics:

- Riemann tensor **singular** on the shock plane
- Null rays & tangent vectors **discontinuous** across shock

Basic properties of a single shock wave II

Symmetries:

- **Axial symmetry** (ϕ_i rotations on $d\Omega_{D-3}$)
- Advanced **null translations** $v \rightarrow v + \text{const.}$
 \Rightarrow Metric of **one shock** wave effectively **2D** (u, ρ)
- **Boosts** along $+z$, with velocity $\beta \equiv \tanh \alpha$, **up to a scaling**

$$E \rightarrow E' = e^\alpha E \quad (\Leftrightarrow \kappa \rightarrow \nu = e^\alpha E)$$

Geometrical optics:

- **Riemann** tensor **singular** on the shock plane
- **Null** rays & tangent vectors **discontinuous** across shock

Basic properties of a single shock wave II

Symmetries:

- **Axial symmetry** (ϕ_i rotations on $d\Omega_{D-3}$)
- Advanced **null translations** $v \rightarrow v + \text{const.}$
 \Rightarrow Metric of **one shock** wave effectively **2D** (u, ρ)
- **Boosts** along $+z$, with velocity $\beta \equiv \tanh \alpha$, **up to a scaling**

$$E \rightarrow E' = e^\alpha E \quad (\Leftrightarrow \kappa \rightarrow \nu = e^\alpha E)$$

Geometrical optics:

- **Riemann** tensor **singular** on the shock plane
- **Null** rays & tangent vectors **discontinuous** across shock

Null rays scattering through shock wave $D = 5$

[Play Video 1](#)

[Play Video 2](#)

- Null rays more **delayed** if aimed **closer to center**
- Also more **bent** inwards, (even backwards) **closer to center**
- Envelope of **outermost null rays defines causal boundary** & becomes original **null plane** of rays @ late times
- Envelope of **innermost rays defines a sphere** @ late times

Null rays scattering through shock wave $D = 4$

[Play Video 1](#)

[Play Video 2](#)

Same as before except that:

- **Rays far from center advance** instead of delaying (coordinate effect since jump $\propto \Phi(\rho)$).
- **But deflection/convergence** of rays produces **same effects** (physical effect since $\propto \frac{d\Phi}{d\rho}$)

1 AS Gravitational Shock Waves

- Physical properties & Ray Optics
- **Superposition & causal structure**
- Perturbative Setup & Solutions


2 Numerical Strategies – Surface Integrals

- Breaking down the problem
- Numerical domain search
- Dealing with the Integrand

Superposition of two shock waves in CM

Consider oppositely moving shocks

⇒ same line element with $u \leftrightarrow v!$



Superposition of two shock waves in CM

Consider oppositely moving shocks


⇒ same line element with $u \leftrightarrow v!$

Superimposing 2 shocks we know:

1 They **move @ speed of light**

⇒ *Cannot influence each other before collision*


⇒ *Metric before collision is sum of two shock metrics*



Superposition of two shock waves in CM

Consider oppositely moving shocks

⇒ same line element with $u \leftrightarrow v!$



Superimposing 2 shocks we know:

1 They **move @ speed of light**

⇒ *Cannot influence each other before collision*

⇒ *Metric before collision is sum of two shock metrics*


2 **How** null generators of a shock **scatter through other shock**

⇒ *Defines causal structure, i.e. regions causally disconnected to collision (see next slide).*

Superposition of two shock waves in CM

Consider oppositely moving shocks

⇒ same line element with $u \leftrightarrow v!$



Superimposing 2 shocks we know:

1 They **move @ speed of light**

⇒ *Cannot influence each other before collision*

⇒ *Metric before collision is sum of two shock metrics*

2 **How** null generators of a shock **scatter through other shock**


⇒ *Defines causal structure, i.e. regions causally disconnected to collision (see next slide).*

3 There is an **apparent horizon** already **before the collision**

⇒ *Black hole must form & Bound on inelasticity is*

$$\epsilon_{\text{radiated}} \leq 1 - \frac{1}{2} \left(\frac{D-2}{2} \frac{\Omega_{D-2}}{\Omega_{D-3}} \right)^{\frac{1}{D-2}}$$

D. M. Eardley and S. B. Giddings, gr-qc/0201034



Causal structure $D = 5$

When summing metrics before collision:

- Easier to interpret metric in **coordinates** (u, v, ρ) adapted to **one shock wave** (e.g. right moving $u = 0$), **since flat**.

Causal structure $D = 5$


When summing metrics before collision:

- Easier to interpret metric in **coordinates** (u, v, ρ) adapted to **one shock wave** (e.g. right moving $u = 0$), **since flat**.
- $ds^2 = ds_{u,flat}^2 + ds_{v,scatter}^2$, where $ds_{v,scatter}^2$ is flat, at points further away from center than envelope of 1st null rays.

Causal structure $D = 5$

When summing metrics before collision:


- Easier to interpret metric in **coordinates** (u, v, ρ) adapted to **one shock wave** (e.g. right moving $u = 0$), **since flat**.
- $ds^2 = ds_{u,flat}^2 + ds_{v,scatter}^2$, where $ds_{v,scatter}^2$ is flat, at points further away from center than envelope of 1st null rays.



Causal structure $D = 5$

When summing metrics before collision:


- Easier to interpret metric in **coordinates** (u, v, ρ) adapted to **one shock wave** (e.g. right moving $u = 0$), **since flat**.
- $ds^2 = ds_{u,flat}^2 + ds_{v,scatter}^2$, where $ds_{v,scatter}^2$ is flat, at points further away from center than envelope of 1st null rays.



Causal structure $D = 5$

When summing metrics before collision:

- Easier to interpret metric in **coordinates** (u, v, ρ) adapted to **one shock wave** (e.g. right moving $u = 0$), **since flat**.
- $ds^2 = ds_{u,flat}^2 + ds_{v,scatter}^2$, where $ds_{v,scatter}^2$ is flat, at points further away from center than envelope of 1st null rays.



Causal structure $D = 4$

Similar diagrams except that:

- Profile function $\lim_{\rho \rightarrow +\infty} \Phi(\rho) = -\infty$.
- Rays with $\rho > \kappa \Rightarrow$ **step forward in time**.

[Play Video 1](#)

[Play Video 2](#)

Causal structure $D = 4$

Similar diagrams except that:

- Profile function $\lim_{\rho \rightarrow +\infty} \Phi(\rho) = -\infty$.
- Rays with $\rho > \kappa \Rightarrow$ **step forward in time**.


[Play Video 1](#)

[Play Video 2](#)

Note (**all** D): backscattered rays ($\frac{\rho_{\text{incident}}}{\kappa^{1/(D-3)}} < 1$) **inside AH** \Rightarrow **Trapped!**

Exact initial conditions in Brinkmann coordinates

In these (asymmetric) coordinates adapted to shock $u = 0$:




Exact initial conditions in Brinkmann coordinates

In these (asymmetric) coordinates adapted to shock $u = 0$:

1 On $u = 0^+$:

$$g_{\mu\nu}(v, x_i) = \eta_{\mu\nu} + \kappa h_{\mu\nu}^{(1)} + \kappa^2 h_{\mu\nu}^{(2)}$$




Exact initial conditions in Brinkmann coordinates

In these (asymmetric) coordinates adapted to shock $u = 0$:

- 1 On $u = 0^+$:

$$g_{\mu\nu}(v, x_i) = \eta_{\mu\nu} + \kappa h_{\mu\nu}^{(1)} + \kappa^2 h_{\mu\nu}^{(2)}$$

- 2 Metric on other surface determined by initial conditions, since **causally connected** only to points on the collision line @ $u = 0$.




Exact initial conditions in Brinkmann coordinates

In these (asymmetric) coordinates adapted to shock $u = 0$:

1 On $u = 0^+$:

$$g_{\mu\nu}(v, x_i) = \eta_{\mu\nu} + \underline{\kappa h_{\mu\nu}^{(1)}} + \underline{\kappa^2 h_{\mu\nu}^{(2)}}$$

2 Metric on other surface determined by initial conditions, since **causally connected** only to points on the collision line @ $u = 0$.



\Rightarrow Well defined evolution problem into (future) **region IV**, given **EXACT** initial conditions.

- 1 AS Gravitational Shock Waves**
 - Physical properties & Ray Optics
 - Superposition & causal structure
 - **Perturbative Setup & Solutions**

- 2 Numerical Strategies – Surface Integrals**
 - Breaking down the problem
 - Numerical domain search
 - Dealing with the Integrand

Two roads to a perturbative construction - Road II

Two roads to a perturbative construction - Road II

Exact initial data on $u = 0$:

- Looks like **series expansion** in κ around Minkowski:

$$g_{\mu\nu}(v, x_i) = \eta_{\mu\nu} + \kappa h_{\mu\nu}^{(1)} + \kappa^2 h_{\mu\nu}^{(2)}$$

Two roads to a perturbative construction - Road II

Exact initial data on $u = 0$:

- Looks like **series expansion** in κ around Minkowski:

$$g_{\mu\nu}(v, x_i) = \eta_{\mu\nu} + \kappa h_{\mu\nu}^{(1)} + \kappa^2 h_{\mu\nu}^{(2)}$$

- In fact $\kappa^i h_{\mu\nu}^{(i)} \propto [\kappa h(v, \rho)]^i \rightarrow [h(v, \rho)]^i$, in units $\kappa = 1$.

Two roads to a perturbative construction - Road II

Exact initial data on $u = 0$:

- Looks like **series expansion** in κ around Minkowski:

$$g_{\mu\nu}(v, x_i) = \eta_{\mu\nu} + \kappa h_{\mu\nu}^{(1)} + \kappa^2 h_{\mu\nu}^{(2)}$$

- In fact $\kappa^i h_{\mu\nu}^{(i)} \propto [\kappa h(v, \rho)]^i \rightarrow [h(v, \rho)]^i$, in units $\kappa = 1$.
- In regions of the (v, ρ) plane where $h(v, \rho) \ll 1$ ($v \ll g(\rho)$):
 \Rightarrow Initial data, 1st and 2nd order perturbation of Minkowski!


Two roads to a perturbative construction - Road II

Exact initial data on $u = 0$:

- Looks like **series expansion** in κ around Minkowski:

$$g_{\mu\nu}(v, x_i) = \eta_{\mu\nu} + \kappa h_{\mu\nu}^{(1)} + \kappa^2 h_{\mu\nu}^{(2)}$$

- In fact $\kappa^i h_{\mu\nu}^{(i)} \propto [\kappa h(v, \rho)]^i \rightarrow [h(v, \rho)]^i$, in units $\kappa = 1$.
- In regions of the (v, ρ) plane where $h(v, \rho) \ll 1$ ($v \ll g(\rho)$):
 \Rightarrow Initial data, 1st and 2nd order perturbation of Minkowski!
- **Condition verified well** for rays of large $\rho_{incident} > 1$.





Two roads to a perturbative construction - Road II

Exact initial data on $u = 0$:

- Looks like **series expansion** in κ around Minkowski:

$$g_{\mu\nu}(v, x_i) = \eta_{\mu\nu} + \kappa h_{\mu\nu}^{(1)} + \kappa^2 h_{\mu\nu}^{(2)}$$

- In fact $\kappa^i h_{\mu\nu}^{(i)} \propto [\kappa h(v, \rho)]^i \rightarrow [h(v, \rho)]^i$, in units $\kappa = 1$.
- In regions of the (v, ρ) plane where $h(v, \rho) \ll 1$ ($v \ll g(\rho)$):
⇒ Initial data, 1st and 2nd order perturbation of Minkowski!
- **Condition verified well** for rays of large $\rho_{incident} > 1$.
⇒ Maps to points close to axis in curved future **region IV**.






Two roads to a perturbative construction - Road II

Exact initial data on $u = 0$:

- Looks like **series expansion** in κ around Minkowski:

$$g_{\mu\nu}(v, x_i) = \eta_{\mu\nu} + \kappa h_{\mu\nu}^{(1)} + \kappa^2 h_{\mu\nu}^{(2)}$$

- In fact $\kappa^i h_{\mu\nu}^{(i)} \propto [\kappa h(v, \rho)]^i \rightarrow [h(v, \rho)]^i$, in units $\kappa = 1$.
- In regions of the (v, ρ) plane where $h(v, \rho) \ll 1$ ($v \ll g(\rho)$):
⇒ Initial data, 1st and 2nd order perturbation of Minkowski!
- **Condition verified well** for rays of large $\rho_{incident} > 1$.
⇒ Maps to points close to axis in curved future **region IV**.



Two roads to a perturbative construction - Road I

D'Eath and Payne's Physical picture:

- 1 Perform a large boost along z with velocity $\beta \equiv \tanh \alpha$
- 2 Energy of $u = 0$ shock $\kappa \rightarrow \nu = e^{\alpha} \kappa$, blue-shifted.
- 3 Energy of $v = 0$ shock $\kappa \rightarrow \lambda = e^{-\alpha} \kappa$, red-shifted.

Two roads to a perturbative construction - Road I

D'Eath and Payne's Physical picture:

- 1 Perform a **large boost along z** with velocity $\beta \equiv \tanh \alpha$
- 2 Energy of $u = 0$ shock $\kappa \rightarrow \nu = e^{\alpha} \kappa$, **blue**-shifted.
- 3 Energy of $v = 0$ shock $\kappa \rightarrow \lambda = e^{-\alpha} \kappa$, **red**-shifted.

Two roads to a perturbative construction - Road I

D'Eath and Payne's Physical picture:

- 1 Perform a **large boost along z** with velocity $\beta \equiv \tanh \alpha$
- 2 Energy of $u = 0$ shock $\kappa \rightarrow \nu = e^{\alpha} \kappa$, **blue**-shifted.
- 3 Energy of $v = 0$ shock $\kappa \rightarrow \lambda = e^{-\alpha} \kappa$, **red**-shifted.

Two roads to a perturbative construction - Road I

D'Eath and Payne's Physical picture:

- 1 Perform a **large boost along z** with velocity $\beta \equiv \tanh \alpha$
- 2 Energy of $u = 0$ shock $\kappa \rightarrow \nu = e^{\alpha} \kappa$, **blue**-shifted.
- 3 Energy of $v = 0$ shock $\kappa \rightarrow \lambda = e^{-\alpha} \kappa$, **red**-shifted.

Two roads to a perturbative construction - Road I

D'Eath and Payne's Physical picture:

- 1 Perform a **large boost along z** with velocity $\beta \equiv \tanh \alpha$
- 2 Energy of $u = 0$ shock $\kappa \rightarrow \nu = e^\alpha \kappa$, **blue**-shifted.
- 3 Energy of $v = 0$ shock $\kappa \rightarrow \lambda = e^{-\alpha} \kappa$, **red**-shifted.

Then:

- **Exact** boundary conditions on $u = 0^+$ (strong shock ν)

$$g_{\mu\nu} = \nu^{\frac{2}{D-3}} \left[\eta_{\mu\nu} + \frac{\lambda}{\nu} h_{\mu\nu}^{(1)} + \left(\frac{\lambda}{\nu} \right)^2 h_{\mu\nu}^{(2)} \right],$$

- Perturbative parameter $\frac{\lambda}{\nu} = e^{-2\alpha} \ll 1$ if α large.

Two roads to a perturbative construction - Road I

D'Eath and Payne's Physical picture:

- 1 Perform a **large boost along z** with velocity $\beta \equiv \tanh \alpha$
- 2 Energy of $u = 0$ shock $\kappa \rightarrow \nu = e^\alpha \kappa$, **blue**-shifted.
- 3 Energy of $v = 0$ shock $\kappa \rightarrow \lambda = e^{-\alpha} \kappa$, **red**-shifted.

Then:

- **Exact** boundary conditions on $u = 0^+$ (strong shock ν)

$$g_{\mu\nu} = \nu^{\frac{2}{D-3}} \left[\eta_{\mu\nu} + \frac{\lambda}{\nu} h_{\mu\nu}^{(1)} + \left(\frac{\lambda}{\nu} \right)^2 h_{\mu\nu}^{(2)} \right],$$

- Perturbative parameter $\frac{\lambda}{\nu} = e^{-2\alpha} \ll 1$ if α large.

Two roads to a perturbative construction - Road I

D'Eath and Payne's Physical picture:

- 1 Perform a **large boost along z** with velocity $\beta \equiv \tanh \alpha$
- 2 Energy of $u = 0$ shock $\kappa \rightarrow \nu = e^\alpha \kappa$, **blue-shifted**.
- 3 Energy of $v = 0$ shock $\kappa \rightarrow \lambda = e^{-\alpha} \kappa$, **red-shifted**.

Then:

- **Exact** boundary conditions on $u = 0^+$ (strong shock ν)

$$g_{\mu\nu} = \nu^{\frac{2}{D-3}} \left[\eta_{\mu\nu} + \frac{\lambda}{\nu} h_{\mu\nu}^{(1)} + \left(\frac{\lambda}{\nu} \right)^2 h_{\mu\nu}^{(2)} \right],$$

- Perturbative parameter $\frac{\lambda}{\nu} = e^{-2\alpha} \ll 1$ if α large.

\Rightarrow Weak shock ($\nu = 0$) small **perturbation of flat space-time.**

Two roads to a perturbative construction - Road I

In Boosted frame

⇒ *Smaller $\rho_{incident}$ allowed.*


⇒ *Perturbative region wider (expands away from the axis).*

Two roads to a perturbative construction - Road I


In Boosted frame

⇒ *Smaller $\rho_{incident}$ allowed.*

⇒ *Perturbative region wider (expands away from the axis).*



CM frame



boosted frame ($-z$ direction)

Boost emphasizes validity of approximation close to the axis.

Perturbative expansion

In the perturbative region (close to rays originating from perturbative initial conditions):

- Assume **perturbative** ansatz

$$g_{\mu\nu}(u > 0, v, x_i) = \eta_{\mu\nu} + \sum_{n=1}^{\infty} \kappa^n h_{\mu\nu}^{(n)}$$

- **Fix gauge** order by order (de Donder condition $\bar{h}^{\alpha\beta}_{,\beta} = 0$)

$$x^\mu \rightarrow x^{N\mu} = x^\mu + \sum_{n=1}^{\infty} \kappa^n \xi^{(n)\mu}$$

- Obtain **decoupled tower of wave equations with source**

$$\square h_{\mu\nu}^{(n)N} = T_{\mu\nu}^{(n-1)} \left[h_{\alpha\beta}^{(k < n)} \right].$$

Note: The source is zero for $i = 1$!

Perturbative expansion

In the perturbative region (close to rays originating from perturbative initial conditions):

- Assume **perturbative** ansatz

$$g_{\mu\nu}(u > 0, v, x_i) = \eta_{\mu\nu} + \sum_{n=1}^{\infty} \kappa^n h_{\mu\nu}^{(n)}$$

- **Fix gauge** order by order (de Donder condition $\bar{h}^{\alpha\beta}_{,\beta} = 0$)

$$x^\mu \rightarrow x^{N\mu} = x^\mu + \sum_{n=1}^{\infty} \kappa^n \xi^{(n)\mu}$$

- Obtain **decoupled tower of wave equations with source**

$$\square h_{\mu\nu}^{(n)N} = T_{\mu\nu}^{(n-1)} \left[h_{\alpha\beta}^{(k < n)} \right].$$

Note: The source is zero for $i = 1$!

Perturbative expansion

In the perturbative region (close to rays originating from perturbative initial conditions):

- Assume **perturbative** ansatz

$$g_{\mu\nu}(u > 0, v, x_i) = \eta_{\mu\nu} + \sum_{n=1}^{\infty} \kappa^n h_{\mu\nu}^{(n)}$$

- **Fix gauge** order by order (de Donder condition $\bar{h}^{\alpha\beta}_{,\beta} = 0$)

$$x^\mu \rightarrow x^{N\mu} = x^\mu + \sum_{n=1}^{\infty} \kappa^n \xi^{(n)\mu}$$

- Obtain **decoupled tower of wave equations with source**

$$\square h_{\mu\nu}^{(n)N} = T_{\mu\nu}^{(n-1)} \left[h_{\alpha\beta}^{(k < n)} \right] .$$

Note: The source is zero for $i = 1$!

Perturbative expansion

In the perturbative region (close to rays originating from perturbative initial conditions):

- Assume **perturbative** ansatz

$$g_{\mu\nu}(u > 0, v, x_i) = \eta_{\mu\nu} + \sum_{n=1}^{\infty} \kappa^n h_{\mu\nu}^{(n)}$$

- **Fix gauge** order by order (de Donder condition $\bar{h}^{\alpha\beta}_{,\beta} = 0$)


$$x^\mu \rightarrow x^{N\mu} = x^\mu + \sum_{n=1}^{\infty} \kappa^n \xi^{(n)\mu}$$

- Obtain **decoupled tower of wave equations with source**

$$\square h_{\mu\nu}^{(n)N} = T_{\mu\nu}^{(n-1)} \left[h_{\alpha\beta}^{(k < n)} \right] .$$

Note: The source is zero for $i = 1$!

Causal structure of the background & formal solution



$$h_{\mu\nu}^{(n)N}(y) = \int_{u'>0} d^D y' G(y, y') \left[T_{\mu\nu}^{(n-1)}(y') + 2\delta(u') \partial_{v'} h_{\mu\nu}^{(n)N}(y') \right]$$

Summary of perturbative solutions I

Axial symmetry allows to expand all metric elements using

$$\Gamma_i \equiv \frac{x_i}{\rho} \quad , \quad \delta_{ij} \quad , \quad \Delta_{ij} \equiv \delta_{ij} - (D-2)\Gamma_i\Gamma_j \quad ,$$

¹ $m = 0, 1, 2$ is the rank of the transverse space-tensor.

Summary of perturbative solutions I

Axial symmetry allows to expand all metric elements using

$$\Gamma_i \equiv \frac{x_i}{\rho}, \quad \delta_{ij}, \quad \Delta_{ij} \equiv \delta_{ij} - (D-2)\Gamma_i\Gamma_j,$$

Then $(A(u, v, \rho), B(u, v, \rho), \dots)$:

$$h_{uu} \equiv A = A^{(1)} + A^{(2)} + \dots \quad h_{ui} \equiv B\Gamma_i = (B^{(1)} + B^{(2)} + \dots)\Gamma_i$$

$$h_{uv} \equiv C = 0 + C^{(2)} + \dots \quad h_{vi} \equiv F\Gamma_i = (0 + F^{(2)} + \dots)\Gamma_i$$

$$h_{vv} \equiv G = 0 + G^{(2)} + \dots$$

$$h_{ij} \equiv E\Delta_{ij} + H\delta_{ij} = (E^{(1)} + E^{(2)} + \dots)\Delta_{ij} + (0 + H^{(2)} + \dots)\delta_{ij}$$

¹ $m = 0, 1, 2$ is the rank of the transverse space-tensor.

Summary of perturbative solutions I

Axial symmetry allows to expand all metric elements using

$$\Gamma_i \equiv \frac{x_i}{\rho}, \quad \delta_{ij}, \quad \Delta_{ij} \equiv \delta_{ij} - (D-2)\Gamma_i\Gamma_j,$$

Then $(A(u, v, \rho), B(u, v, \rho), \dots)$:

$$h_{uu} \equiv A = A^{(1)} + A^{(2)} + \dots \quad h_{ui} \equiv B\Gamma_i = (B^{(1)} + B^{(2)} + \dots)\Gamma_i$$

$$h_{uv} \equiv C = 0 + C^{(2)} + \dots \quad h_{vi} \equiv F\Gamma_i = (0 + F^{(2)} + \dots)\Gamma_i$$

$$h_{vv} \equiv G = 0 + G^{(2)} + \dots$$

$$h_{ij} \equiv E\Delta_{ij} + H\delta_{ij} = (E^{(1)} + E^{(2)} + \dots)\Delta_{ij} + (0 + H^{(2)} + \dots)\delta_{ij}$$

Similar decomposition for the source $T_{\mu\nu}^{(i-1)}$. Also denote:

- Generic metric function by $F_m^{(n)}(u, v, \rho)^1$.
- Its source by $S_F^{(n-1)}(u, v, \rho)$.
- Note $F_m^{(n)}(0, v, \rho) = f(\rho) [h(v, \rho)]^n$, $n = 1, 2$.

¹ $m = 0, 1, 2$ is the rank of the transverse space-tensor.

Summary of perturbative solutions I

Axial symmetry allows to expand all metric elements using

$$\Gamma_i \equiv \frac{x_i}{\rho}, \quad \delta_{ij}, \quad \Delta_{ij} \equiv \delta_{ij} - (D-2)\Gamma_i\Gamma_j,$$

Then $(A(u, v, \rho), B(u, v, \rho), \dots)$:

$$h_{uu} \equiv A = A^{(1)} + A^{(2)} + \dots \quad h_{ui} \equiv B\Gamma_i = (B^{(1)} + B^{(2)} + \dots)\Gamma_i$$

$$h_{uv} \equiv C = 0 + C^{(2)} + \dots \quad h_{vi} \equiv F\Gamma_i = (0 + F^{(2)} + \dots)\Gamma_i$$

$$h_{vv} \equiv G = 0 + G^{(2)} + \dots$$

$$h_{ij} \equiv E\Delta_{ij} + H\delta_{ij} = (E^{(1)} + E^{(2)} + \dots)\Delta_{ij} + (0 + H^{(2)} + \dots)\delta_{ij}$$

Similar decomposition for the source $T_{\mu\nu}^{(i-1)}$. Also denote:

■ Generic metric function by $F_m^{(n)}(u, v, \rho)^1$.

■ Its source by $S_F^{(n-1)}(u, v, \rho)$.

■ Note $F_m^{(n)}(0, v, \rho) = f(\rho)[h(v, \rho)]^n$, $n = 1, 2$.

¹ $m = 0, 1, 2$ is the rank of the transverse space-tensor.

Summary of perturbative solutions I

Axial symmetry allows to expand all metric elements using

$$\Gamma_i \equiv \frac{x_i}{\rho}, \quad \delta_{ij}, \quad \Delta_{ij} \equiv \delta_{ij} - (D-2)\Gamma_i\Gamma_j,$$

Then $(A(u, v, \rho), B(u, v, \rho), \dots)$:

$$h_{uu} \equiv A = A^{(1)} + A^{(2)} + \dots \quad h_{ui} \equiv B\Gamma_i = (B^{(1)} + B^{(2)} + \dots)\Gamma_i$$

$$h_{uv} \equiv C = 0 + C^{(2)} + \dots \quad h_{vi} \equiv F\Gamma_i = (0 + F^{(2)} + \dots)\Gamma_i$$

$$h_{vv} \equiv G = 0 + G^{(2)} + \dots$$

$$h_{ij} \equiv E\Delta_{ij} + H\delta_{ij} = (E^{(1)} + E^{(2)} + \dots)\Delta_{ij} + (0 + H^{(2)} + \dots)\delta_{ij}$$

Similar decomposition for the source $T_{\mu\nu}^{(i-1)}$. Also denote:

■ Generic metric function by $F_m^{(n)}(u, v, \rho)^1$.

■ Its source by $S_F^{(n-1)}(u, v, \rho)$.

■ Note $F_m^{(n)}(0, v, \rho) = f(\rho)[h(v, \rho)]^n$, $n = 1, 2$.

¹ $m = 0, 1, 2$ is the rank of the transverse space-tensor.

Summary of perturbative solutions I

Axial symmetry allows to expand all metric elements using

$$\Gamma_i \equiv \frac{x_i}{\rho}, \quad \delta_{ij}, \quad \Delta_{ij} \equiv \delta_{ij} - (D-2)\Gamma_i\Gamma_j,$$

Then $(A(u, v, \rho), B(u, v, \rho), \dots)$:

$$h_{uu} \equiv A = A^{(1)} + A^{(2)} + \dots \quad h_{ui} \equiv B\Gamma_i = (B^{(1)} + B^{(2)} + \dots)\Gamma_i$$

$$h_{uv} \equiv C = 0 + C^{(2)} + \dots \quad h_{vi} \equiv F\Gamma_i = (0 + F^{(2)} + \dots)\Gamma_i$$

$$h_{vv} \equiv G = 0 + G^{(2)} + \dots$$

$$h_{ij} \equiv E\Delta_{ij} + H\delta_{ij} = (E^{(1)} + E^{(2)} + \dots)\Delta_{ij} + (0 + H^{(2)} + \dots)\delta_{ij}$$

Similar decomposition for the source $T_{\mu\nu}^{(i-1)}$. Also denote:

- Generic metric function by $F_m^{(n)}(u, v, \rho)^1$.
- Its source by $S_F^{(n-1)}(u, v, \rho)$.
- Note $F_m^{(n)}(0, v, \rho) = f(\rho)[h(v, \rho)]^n$, $n = 1, 2$.

¹ $m = 0, 1, 2$ is the rank of the transverse space-tensor.

Summary of perturbative solutions II

Then the integral solution is:

$$F_m^{(n)}(u, v, \rho) = F_{m, \text{Surf}}^{(n)} + F_{m, \text{Vol}}^{(n)}$$

Summary of perturbative solutions II

Then the integral solution is:

$$F_m^{(n)}(u, v, \rho) = F_{m, \text{Surf}}^{(n)} + F_{m, \text{Vol}}^{(n)}$$

with

$$F_{m, \text{Surf}}^{(n)} = -\frac{n!(-1)^D \Omega_{D-4}}{(2\pi\rho)^{\frac{D-2}{2}}} \left(\frac{\sqrt{2}\rho}{u}\right)^n \int_{\mathcal{D}_{\text{Surf}}} d\rho' f(\rho') \rho'^{\frac{D-4}{2}+n} I_m^{D,n}(x_\star)$$

Summary of perturbative solutions II

Then the integral solution is:

$$F_m^{(n)}(u, v, \rho) = F_{m, \text{Surf}}^{(n)} + F_{m, \text{Vol}}^{(n)}$$

with

$$F_{m, \text{Surf}}^{(n)} = -\frac{n!(-1)^D \Omega_{D-4}}{(2\pi\rho)^{\frac{D-2}{2}}} \left(\frac{\sqrt{2}\rho}{u}\right)^n \int_{\mathcal{D}_{\text{Surf}}} d\rho' f(\rho') \rho'^{\frac{D-4}{2}+n} I_m^{D,n}(x_*)$$

and

$$F_{m, \text{Vol}}^{(n)} = \frac{-\Omega_{D-4}}{2(2\pi\rho)^{\frac{D-2}{2}}} \int_0^u du' \iint_{\mathcal{D}_{\text{Vol}}} dv' d\rho' \rho'^{\frac{D-4}{2}} S_F^{(n-1)}(u', v', \rho') I_m^{D,0}(x)$$

Summary of perturbative solutions II

Then the integral solution is:

$$F_m^{(n)}(u, v, \rho) = F_{m, \text{Surf}}^{(n)} + F_{m, \text{Vol}}^{(n)}$$

with

$$F_{m, \text{Surf}}^{(n)} = -\frac{n!(-1)^D \Omega_{D-4}}{(2\pi\rho)^{\frac{D-2}{2}}} \left(\frac{\sqrt{2}\rho}{u}\right)^n \int_{\mathcal{D}_{\text{Surf}}} d\rho' f(\rho') \rho'^{\frac{D-4}{2}+n} I_m^{D,n}(x_*)$$


and

$$F_{m, \text{Vol}}^{(n)} = \frac{-\Omega_{D-4}}{2(2\pi\rho)^{\frac{D-2}{2}}} \int_0^u du' \iint_{\mathcal{D}_{\text{Vol}}} dv' d\rho' \rho'^{\frac{D-4}{2}} S_F^{(n-1)}(u', v', \rho') I_m^{D,0}(x)$$

$$x = \frac{\rho^2 + \rho'^2 - 2(u-u')(v-v')}{2\rho\rho'} \leq 1$$

$x_* = x|_{\text{@ collision line}}$

($I_m^{D,n}$ in handout).



Summary of perturbative solutions II

Then the integral solution is:

$$F_m^{(n)}(u, v, \rho) = F_{m, \text{Surf}}^{(n)} + F_{m, \text{Vol}}^{(n)}$$

with

$$F_{m, \text{Surf}}^{(n)} = -\frac{n!(-1)^D \Omega_{D-4}}{(2\pi\rho)^{\frac{D-2}{2}}} \left(\frac{\sqrt{2}\rho}{u}\right)^n \int_{\mathcal{D}_{\text{Surf}}} d\rho' f(\rho') \rho'^{\frac{D-4}{2}+n} I_m^{D,n}(x_*)$$


and

$$F_{m, \text{Vol}}^{(n)} = \frac{-\Omega_{D-4}}{2(2\pi\rho)^{\frac{D-2}{2}}} \int_0^u du' \iint_{\mathcal{D}_{\text{Vol}}} dv' d\rho' \rho'^{\frac{D-4}{2}} S_F^{(n-1)}(u', v', \rho') I_m^{D,0}(x)$$

$$x = \frac{\rho^2 + \rho'^2 - 2(u-u')(v-v')}{2\rho\rho'} \leq 1$$

x_* = x | @ collision line

($I_m^{D,n}$ in handout).



Physical interpretation of the solutions

A very clear physical picture to this tower of solutions is:

- 1 At 1st order, there is an impulsive creation of a configuration of radiation on $u = 0^+$ in the initial conditions. This propagates freely on the flat background.
- 2 The 2nd order $h_{\mu\nu}^{(2)}$ propagates on a flat background, and another signal of radiation is generated by the source of $h_{\mu\nu}^{(1)}$. This must encode backscattering.
- 3 Higher order perturbations $h_{\mu\nu}^{(i>2)}$, are zero on $u = 0$. However, they are sourced by lower order perturbations, thus radiation is created spontaneously.

Physical interpretation of the solutions

A very clear physical picture to this tower of solutions is:

- 1 At 1st order, there is an impulsive creation of a configuration of radiation on $u = 0^+$ in the initial conditions. This propagates freely on the flat background.
- 2 The 2nd order $h_{\mu\nu}^{(2)}$ propagates on a flat background, and another signal of radiation is generated by the source of $h_{\mu\nu}^{(1)}$. This must encode backscattering.
- 3 Higher order perturbations $h_{\mu\nu}^{(i>2)}$, are zero on $u = 0$. However, they are sourced by lower order perturbations, thus radiation is created spontaneously.

Physical interpretation of the solutions

A very clear physical picture to this tower of solutions is:

- 1 At 1st order, there is an impulsive creation of a configuration of radiation on $u = 0^+$ in the initial conditions. This propagates freely on the flat background.
- 2 The 2nd order $h_{\mu\nu}^{(2)}$ propagates on a flat background, and another signal of radiation is generated by the source of $h_{\mu\nu}^{(1)}$. This must encode backscattering.
- 3 Higher order perturbations $h_{\mu\nu}^{(i>2)}$, are zero on $u = 0$. However, they are sourced by lower order perturbations, thus radiation is created spontaneously.

Physical interpretation of the solutions

A very clear physical picture to this tower of solutions is:

- 1 At 1st order, there is an impulsive creation of a configuration of radiation on $u = 0^+$ in the initial conditions. This propagates freely on the flat background.
- 2 The 2nd order $h_{\mu\nu}^{(2)}$ propagates on a flat background, and another signal of radiation is generated by the source of $h_{\mu\nu}^{(1)}$. This must encode backscattering.
- 3 Higher order perturbations $h_{\mu\nu}^{(i>2)}$, are zero on $u = 0$. However, they are sourced by lower order perturbations, thus radiation is created spontaneously.

- 1 AS Gravitational Shock Waves
 - Physical properties & Ray Optics
 - Superposition & causal structure
 - Perturbative Setup & Solutions

- 2 Numerical Strategies – Surface Integrals
 - Breaking down the problem
 - Numerical domain search
 - Dealing with the Integrand

Surface integrals & Simplified form

Change variables $y \equiv \frac{\rho'}{\rho}$, $\rho \equiv (\sqrt{2}v - \Phi(\rho))\rho^{D-4}$ and $q \equiv \frac{u}{\rho^{D-2}}$:

Surface integrals & Simplified form

Change variables $y \equiv \frac{\rho'}{\rho}$, $\rho \equiv (\sqrt{2}v - \Phi(\rho))\rho^{D-4}$ and $q \equiv \frac{u}{\rho^{D-2}}$:

$$F_{m, \text{Surf}}^{(n)} = - \frac{n!(-1)^D \Omega_{D-4}}{\rho^{(D-3)(2n+N_u-N_v)} (2\pi)^{\frac{D-2}{2}}} \left(\frac{\sqrt{2}}{q}\right)^n \int_{\mathcal{D}_{\text{Surf}}} dy f(y) y^{\frac{D-4}{2} + n} I_m^{D,n}(x_*)$$

Surface integrals & Simplified form

Change variables $y \equiv \frac{\rho'}{\rho}$, $\rho \equiv (\sqrt{2}v - \Phi(\rho))\rho^{D-4}$ and $q \equiv \frac{u}{\rho^{D-2}}$:

$$F_{m, \text{Surf}}^{(n)} = - \frac{n!(-1)^D \Omega_{D-4}}{\rho^{(D-3)(2n+N_u-N_v)} (2\pi)^{\frac{D-2}{2}}} \left(\frac{\sqrt{2}}{q}\right)^n \int_{\mathcal{D}_{\text{Surf}}} dy f(y) y^{\frac{D-4}{2} + n} I_m^{D,n}(x_*)$$

$$x_* = \frac{1 + y^2 - \sqrt{2}q(\rho - \psi(y))}{2y}, \quad \psi(y) \equiv \begin{cases} -2 \log y & , D = 4 \\ \frac{2}{D-4} \left[\frac{1}{y^{D-4}} - 1 \right] & , D > 4 \end{cases}$$

- ρ dependence scales out!
- Integrand depends only on p, q (hidden symmetry)!
- Non-trivial domain $\mathcal{D}_{\text{Surf}} : x_* \leq 1$.

Surface integrals & Simplified form

Change variables $y \equiv \frac{\rho'}{\rho}$, $\rho \equiv (\sqrt{2}v - \Phi(\rho))\rho^{D-4}$ and $q \equiv \frac{u}{\rho^{D-2}}$:

$$F_{m, \text{Surf}}^{(n)} = - \frac{n!(-1)^D \Omega_{D-4}}{\rho^{(D-3)(2n+N_u-N_v)} (2\pi)^{\frac{D-2}{2}}} \left(\frac{\sqrt{2}}{q}\right)^n \int_{\mathcal{D}_{\text{Surf}}} dy f(y) y^{\frac{D-4}{2} + n} I_m^{D,n}(\mathbf{x}_\star)$$

$$\mathbf{x}_\star = \frac{1 + y^2 - \sqrt{2}q(\rho - \psi(y))}{2y}, \quad \psi(y) \equiv \begin{cases} -2 \log y & , D = 4 \\ \frac{2}{D-4} \left[\frac{1}{y^{D-4}} - 1 \right] & , D > 4 \end{cases}$$

- ρ dependence scales out!
- Integrand depends only on p, q (hidden symmetry)!
- Non-trivial domain $\mathcal{D}_{\text{Surf}} : \mathbf{x}_\star \leq 1$.

Summary of the procedure

The numerical task at hand then involves the following steps:

Summary of the procedure

The numerical task at hand then involves the following steps:

- 1** *Numerical search of integration domain $\mathcal{D}_{Surf}, x_* < 1$:
Classification of **roots & 1D bracketing**.*
- 2** *Implementation of functions $I_m^{D,n}(x_*)$: Polynomial (or series) **expansions & recurrence relations**.*
- 3** *Removal of integrable singularities of $I_m^{D,n}(x_*)$: adapted **changes of variables & polynomial (or series) expansions**.*

Summary of the procedure

The numerical task at hand then involves the following steps:

- 1** *Numerical search of integration domain \mathcal{D}_{Surf} , $x_* < 1$:*
Classification of **roots & 1D bracketing**.
- 2** *Implementation of functions $I_m^{D,n}(x_*)$:* Polynomial (or series) **expansions & recurrence relations**.
- 3** *Removal of integrable singularities of $I_m^{D,n}(x_*)$:* adapted **changes of variables** & polynomial (or series) **expansions**.

Summary of the procedure

The numerical task at hand then involves the following steps:

- 1** *Numerical search of integration domain \mathcal{D}_{Surf} , $x_* < 1$:*
Classification of **roots & 1D bracketing**.
- 2** *Implementation of functions $I_m^{D,n}(x_*)$:* Polynomial (or series) **expansions & recurrence relations**.
- 3** *Removal of integrable singularities of $I_m^{D,n}(x_*)$:* adapted **changes of variables** & polynomial (or series) **expansions**.

Summary of the procedure

The numerical task at hand then involves the following steps:

- 1** *Numerical search of integration domain $\mathcal{D}_{Surf}, x_* < 1$:*
Classification of **roots & 1D bracketing**.
- 2** *Implementation of functions $I_m^{D,n}(x_*)$:* Polynomial (or series) **expansions & recurrence relations**.
- 3** *Removal of integrable singularities of $I_m^{D,n}(x_*)$:* adapted **changes of variables** & polynomial (or series) **expansions**.

Main code written in C++ with GSL libraries for numerical integration & bracketing algorithms.

I will show Mathematica checks/examples to illustrate.

General advice from my experiences with numerics

General advice from my experiences with numerics



Break down the problem (self contained/testable tasks)

General advice from my experiences with numerics



Break down the problem (self contained/testable tasks)

Test your code to destruction: The earlier the better!



General advice from my experiences with numerics



Break down the problem (self contained/testable tasks)

Test your code to destruction: The earlier the better!



Be careful how you write expressions!

General advice from my experiences with numerics



Break down the problem (self contained/testable tasks)

Test your code to destruction: The earlier the better!



Be careful how you write expressions!

Identify operations to be done once instead of repeatedly



General advice from my experiences with numerics



Break down the problem (self contained/testable tasks)

Test your code to destruction: The earlier the better!



Be careful how you write expressions!

Identify operations to be done once instead of repeatedly



Be patient, take your time to code. It saves you time!

- 1 AS Gravitational Shock Waves
 - Physical properties & Ray Optics
 - Superposition & causal structure
 - Perturbative Setup & Solutions

- 2 Numerical Strategies – Surface Integrals
 - Breaking down the problem
 - Numerical domain search
 - Dealing with the Integrand

Domain conditions

Consider the two following polynomials/functions ($s = \pm$)

$$C_s(y) \equiv y^{D-2} + 2sy^{D-3} - (\sqrt{2}qp - 1)y^{D-4} + \sqrt{2}qy^{D-4}\psi(y)$$

Domain conditions

Consider the two following polynomials/functions ($s = \pm$)

$$C_s(y) \equiv y^{D-2} + 2sy^{D-3} - (\sqrt{2}qp - 1)y^{D-4} + \sqrt{2}qy^{D-4}\psi(y)$$

Past light cone/collision condition $x_* \leq 1$ reduces to $C_-(y) \leq 0$.

See later $x_* = -1$ induces integrable **singularity @ $C_+(y) = 0$!**

Domain conditions

Consider the two following polynomials/functions ($s = \pm$)

$$C_s(y) \equiv y^{D-2} + 2sy^{D-3} - (\sqrt{2}qp - 1)y^{D-4} + \sqrt{2}qy^{D-4}\psi(y)$$

Past light cone/collision condition $x_* \leq 1$ reduces to $C_-(y) \leq 0$.

See later $x_* = -1$ induces integrable **singularity @ $C_+(y) = 0$!**

One can also show that:

- $C_+(y) \geq C_-(y)$.
- For $y \geq 0$ there are ≤ 2 roots for $C_s(y)$ & a minimum at

$$y_{min} = \frac{-s(D-3) + \sqrt{1 + \sqrt{2}q(D-2)((D-4)p+2)}}{D-2}$$

Domain conditions

Consider the two following polynomials/functions ($s = \pm$)

$$C_s(y) \equiv y^{D-2} + 2sy^{D-3} - (\sqrt{2}qp - 1)y^{D-4} + \sqrt{2}qy^{D-4}\psi(y)$$

Past light cone/collision condition $x_* \leq 1$ reduces to $C_-(y) \leq 0$.

See later $x_* = -1$ induces integrable **singularity @ $C_+(y) = 0$!**

One can also show that:

- $C_+(y) \geq C_-(y)$.
- For $y \geq 0$ there are **≤ 2 roots** for $C_s(y)$ & a **minimum at**

$$y_{min} = \frac{-s(D-3) + \sqrt{1 + \sqrt{2}q(D-2)((D-4)p+2)}}{D-2}$$

Domain conditions

Consider the two following polynomials/functions ($s = \pm$)

$$C_s(y) \equiv y^{D-2} + 2sy^{D-3} - (\sqrt{2}qp - 1)y^{D-4} + \sqrt{2}qy^{D-4}\psi(y)$$

Past light cone/collision condition $x_* \leq 1$ reduces to $C_-(y) \leq 0$.

See later $x_* = -1$ induces integrable **singularity @ $C_+(y) = 0$!**

One can also show that:

- $C_+(y) \geq C_-(y)$.
- For $y \geq 0$ there are **≤ 2 roots** for $C_s(y)$ & a **minimum at**

$$y_{min} = \frac{-s(D-3) + \sqrt{1 + \sqrt{2}q(D-2)((D-4)p+2)}}{D-2}$$

\Rightarrow Domain determination reduces **to root finding**
(see Mathematica example 1)

- 1 AS Gravitational Shock Waves
 - Physical properties & Ray Optics
 - Superposition & causal structure
 - Perturbative Setup & Solutions

- 2 Numerical Strategies – Surface Integrals
 - Breaking down the problem
 - Numerical domain search
 - Dealing with the Integrand

Sensitive features of the integrand $I_m^{D,n}(x_*)$

Integrand contains several pieces **to be computed with care**:

- 1 **Polynomials** $Q_m^{D,n}(x_*)$, ... coefficients defined differentially:

Example of task to be done once at the beginning.

(see Mathematica example 2.1)

- 2 D odd contains a potentially **problematic scaling**.

Example where asymptotic expansions can be useful!

(see Mathematica example 2.2)

- 3 **Singularities** @ $x_* = s = \pm 1$, due to

$$\frac{1}{\sqrt{\pm(1-x_*^2)}} = \frac{1}{\sqrt{(1-x_*)(\pm x_* \pm 1)}}$$

- Adapted change of variable $y = y_s^{\text{root}} - kw^2$ & split domains.
- Writing expressions & using series in problematic regions.

(see Mathematica example 2.3)

Sensitive features of the integrand $I_m^{D,n}(x_*)$

Integrand contains several pieces **to be computed with care**:

1 **Polynomials** $Q_m^{D,n}(x_*)$, ... coefficients defined differentially:
Example of task to be done once at the beginning.
(see Mathematica example 2.1)

2 D odd contains a potentially **problematic scaling**.
Example where asymptotic expansions can be useful!
(see Mathematica example 2.2)

3 **Singularities** @ $x_* = s = \pm 1$, due to

$$\frac{1}{\sqrt{\pm(1-x_*^2)}} = \frac{1}{\sqrt{(1-x_*)(\pm x_* \pm 1)}}$$

- Adapted change of variable $y = y_s^{\text{root}} - kw^2$ & split domains.
- Writing expressions & using series in problematic regions.

(see Mathematica example 2.3)

Sensitive features of the integrand $I_m^{D,n}(x_*)$

Integrand contains several pieces **to be computed with care**:

1 Polynomials $Q_m^{D,n}(x_*)$, ... coefficients defined differentially:
Example of task to be done once at the beginning.
(see Mathematica example 2.1)

2 D odd contains a potentially **problematic scaling**.
Example where asymptotic expansions can be useful!
(see Mathematica example 2.2)

3 Singularities @ $x_* = s = \pm 1$, due to

$$\frac{1}{\sqrt{\pm(1-x_*^2)}} = \frac{1}{\sqrt{(1-x_*)(\pm x_* \pm 1)}}$$

- Adapted change of variable $y = y_s^{\text{root}} - kw^2$ & split domains.
- Writing expressions & using series in problematic regions.

(see Mathematica example 2.3)

Sensitive features of the integrand $I_m^{D,n}(x_*)$

Integrand contains several pieces **to be computed with care**:

1 Polynomials $Q_m^{D,n}(x_*)$, ... coefficients defined differentially:
Example of task to be done once at the beginning.
(see Mathematica example 2.1)

2 D odd contains a potentially **problematic scaling**.
Example where asymptotic expansions can be useful!
(see Mathematica example 2.2)

3 **Singularities** @ $x_* = s = \pm 1$, due to

$$\frac{1}{\sqrt{\pm(1-x_*^2)}} = \frac{1}{\sqrt{(1-x_*)(\pm x_* \pm 1)}}$$

- Adapted change of variable $y = y_s^{\text{root}} - kw^2$ & split domains.
- Writing expressions & using series in problematic regions.

(see Mathematica example 2.3)

So far we have seen:

- How to **superimpose** two shock waves & write exact initial conditions for evolution;
- Formulated approximate **perturbative scheme** to find gravitational perturbations close to axis;
- Found the general perturbative solutions, order by order, as **source and volume integrals**;
- Addressed surface integrals with some **practical examples**.

Tomorrow \Rightarrow Numerical results, radiation extraction
& Higher orders.